
Recombinant X-Prolyl Aminopeptidase 3 (XPNPEP3)(N-6His)

Catalog#:PEH25156

概述（Summary）

别称（Alternative Names）	APP3; NPHPL1; Aminopeptidase P3
纯度（Purity）	Greater than 90% as determined by reducing SDS-PAGE.
内毒素（Endotoxin level）	<1 EU/μg as determined by LAL test.
蛋白构建（Construction）	6His-【Glu253-Gln482】
Accession #	Q9NQH7
表达宿主（Host）	E.coli
种属（Species）	Homo sapiens (Human)
预测分子量（Predicted MW）	29.4 kDa
制剂（Form）	Lyophilized from a 0.2 μm filtered solution of 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.
运输方式（Shipping）	The product is shipped with ice bag. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存（Stability &Storage）	12 months at -20°C to -80°C in lyophilized state.6 months at -20°C to -80°C under sterile conditions after reconstitution.7-10 days at 2°C to 8°C under sterile conditions after reconstitution.Please minimize freeze-thaw cycles.
复溶（Reconstitution）	Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.
背景介绍（References）	Catalyzes the removal of a penultimate prolyl residue from the N-termini of peptides, such as Leu-Pro-Ala. Also shows low activity towards peptides with Ala or Ser at the P1 position.; Promotes TNFRSF1B-mediated phosphorylation of MAPK8/JNK1 and MAPK9/JNK2, suggesting a function as an adapter protein for TNFRSF1B; the effect is independent of XPNPEP3 peptidase activity. May inhibit apoptotic cell death induced via TNF-TNFRSF1B signaling.

电泳图（SDS-PAGE image）



Greater than 90% as determined by reducing SDS-PAGE.

注意事项（Note）

For Research Use Only , Not for Diagnostic Use.