
Active Granzyme K (GZMK)

Catalog#:PEM12623

概述（Summary）

别称（Alternative Names）	GZM-K; TRYP2; PRSS; Serine Protease,Granzyme 3; Tryptase II; Fragmentin-3; Granzyme-3; NK-tryptase-2
纯度（Purity）	Greater than 90% as determined by reducing SDS-PAGE.
内毒素（Endotoxin level）	Please contact with the lab for this information
蛋白构建（Construction）	【Gln44-Val227】
Accession #	O35205
表达宿主（Host）	E.coli
种属（Species）	Mus musculus (Mouse)
预测分子量（Predicted MW）	24.1 kDa
制剂（Form）	Lyophilized from a 0.2 μm filtered solution of PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and Proclin300.
运输方式（Shipping）	The product is shipped with ice bag. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存（Stability &Storage）	12 months at -20°C to -80°C in lyophilized state.6 months at -20°C to -80°C under sterile conditions after reconstitution.7-10 days at 2°C to 8°C under sterile conditions after reconstitution.Please minimize freeze-thaw cycles.
复溶（Reconstitution）	Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

电泳图（SDS-PAGE image）



Greater than 90% as determined by reducing SDS-PAGE.

活性图（Bioactivity image）

$$\text{Specific Activity (pmol/min/ug)} = \frac{\text{Adjusted } V_{max}^{*} \text{ (OD/min) x well volume (L) x } 10^{12} \text{ pmol/mol}}{\text{ext. coeff}^{**} \text{ } (M^{-1}cm^{-1}) \text{ x path corr.}^{***} \text{ (cm) x amount of enzyme (ug)}}$$

*Adjusted for Substrate Blank
**Using the extinction coefficient 13800 $M^{-1}cm^{-1}$
***Using the path correction 0.320 cm

Granzyme K is a member of the granzyme family of the serine proteases found specifically in the cytotoxic granules of cytotoxic T lymphocytes (CTL) and natural killer (NK) cells. Mouse granzyme K is synthesized as a precursor (263 residues) with a signal peptide (residues 1 21), a propeptide (residues 22-25) and a mature chain (residues 26-263). The activity of recombinant mouse Granzyme K was measured by its ability to cleaves a thioester substrate Z-Lys-SBzl•HCl. The reaction was performed in 0.05 M Tris, 0.15 M NaCl, 0.01% Triton X-100, pH 8.0 (assay buffer), initiated by addition 50 µL of various concentrations of GZMK (diluted by assay buffer) to 50 µL of 1.2 mM substrate and DTNB mixture. The final well serves as a negative control with no GZMK, replace with 50 µL assay buffer. Incubated at 25℃ for 5min, then read at a wavelength of 405 nm. The specific activity of recombinant mouse Granzyme K is >70 pmol/min/µg.

注意事项（Note）

For Research Use Only , Not for Diagnostic Use.