
# Recombinant Glycoprotein 2, Zymogen Granule Membrane (GP2)(N-6His)

*Catalog#:PEM22127*

## 概述（Summary）

| | |
|---|---|
| **别称（Alternative Names）** | ZAP75; Pancreatic Secretory Granule Membrane Major Glycoprotein GP2; Pancreatic zymogen granule membrane protein GP-2 |
| **纯度（Purity）** | Greater than 97% as determined by reducing SDS-PAGE. |
| **内毒素（Endotoxin level）** | <1 EU/μg as determined by LAL test. |
| **蛋白构建（Construction）** | 6His-【Phe301-Lys506】 |
| **Accession #** | Q9D733 |
| **表达宿主（Host）** | E.coli |
| **种属（Species）** | Mus musculus (Mouse) |
| **预测分子量（Predicted MW）** | 26.9 kDa |
| **制剂（Form）** | Lyophilized from a 0.2 μm filtered solution of PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and Proclin300. |
| **运输方式（Shipping）** | The product is shipped with ice bag. Upon receipt, store it immediately at the temperature listed below. |
| **稳定性&储存（Stability &Storage）** | 12 months at -20°C to -80°C in lyophilized state.6 months at -20°C to -80°C under sterile conditions after reconstitution.7-10 days at 2°C to 8°C under sterile conditions after reconstitution.Please minimize freeze-thaw cycles. |
| **复溶（Reconstitution）** | Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex. |
| **背景介绍（References）** | Functions as an intestinal M-cells transcytotic receptor specific of type-I-piliated bacteria that participate in the mucosal immune response toward these bacteria. At the apical membrane of M-cells binds fimH, a protein of the bacteria type I pilus tip. Internalizes bound bacteria, like E.coli and S.typhimurium, from the lumen of the intestine and delivers them, through M-cells, to the underlying organized lymphoid follicles where they are captured by antigen-presenting dendritic cells to ellicit a mucosal immune response. |

## 电泳图（SDS-PAGE image）



15% SDS-PAGE

Greater than 97% as determined by reducing SDS-PAGE.

## 注意事项（Note）

For Research Use Only , Not for Diagnostic Use.