Recombinant Death Associated Protein Kinase 3 (DAPK3)(N-6His)


Catalog#:PEM22577

概述（Summary）

别称（Alternative Names）	ZIP; ZIPK; Dlk; DAP-like kinase; MYPT1 kinase; Zipper-interacting protein kinase
纯度（Purity）	Greater than 97% as determined by reducing SDS-PAGE.
内毒素（Endotoxin level）	<1 EU/μg as determined by LAL test.
蛋白构建（Construction）	6His-【Met1-Arg448】
Accession #	O54784
表达宿主（Host）	E.coli
种属（Species）	Mus musculus (Mouse)
预测分子量（Predicted MW）	55.1 kDa
制剂（Form）	Lyophilized from a 0.2 μm filtered solution of PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and Proclin300.
运输方式（Shipping）	The product is shipped with ice bag. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存（Stability &Storage）	12 months at -20°C to -80°C in lyophilized state.6 months at -20°C to -80°C under sterile conditions after reconstitution.7-10 days at 2°C to 8°C under sterile conditions after reconstitution.Please minimize freeze-thaw cycles.
复溶（Reconstitution）	Reconstitute in ddH_2O to a concentration of 0.1-0.2 mg/mL. Do not vortex.
背景介绍（References）	Serine/threonine kinase which is involved in the regulation of apoptosis, autophagy, transcription, translation and actin cytoskeleton reorganization. Regulates both type I (caspase-dependent) apoptotic and type II (caspase-independent) autophagic cell deaths signal, depending on the cellular setting. Involved in formation of promyelocytic leukemia protein nuclear body (PML-NB). Involved in apoptosis involving PAWR which mediates cytoplasmic relocation; in vitro phosphorylates PAWR. Phosphorylates MYL12B in non-muscle cells leading to reorganization of actin cytoskeleton such as in regulation of cell polarity and cell migration. Positively regulates canonical Wnt/beta-catenin signaling through interaction with NLK and TCF7L2; disrupts the NLK-TCF7L2 complex thereby

influencing the phosphorylation of TCF7L2 by NLK. Phosphorylates STAT3 and enhances its transcriptional activity. Enhances transcription from AR-responsive promoters in a hormone- and kinase-dependent manner. Phosphorylates histone H3 on 'Thr-11' at centromeres during mitosis. Phosphorylates RPL13A on 'Ser-77' upon interferon-gamma activation which is causing RPL13A release from the ribosome, RPL13A association with the GAIT complex and its subsequent involvement in transcript-selective translation inhibition.

电泳图（SDS-PAGE image）



Greater than 97% as determined by reducing SDS-PAGE.

注意事项（Note）

For Research Use Only , Not for Diagnostic Use.