
# Active Ephrin A4 (EFNA4)

*Catalog#:PER12926*

## 概述（Summary）

| | |
|---|---|
| **别称（Alternative Names）** | EFL4; EPLG4; LERK4; EPH-related receptor tyrosine kinase ligand 4 |
| **纯度（Purity）** | Greater than 90% as determined by reducing SDS-PAGE. |
| **内毒素（Endotoxin level）** | Please contact with the lab for this information |
| **蛋白构建（Construction）** | 【Met1-Leu204】 |
| **Accession #** | D3ZEN1 |
| **表达宿主（Host）** | E.coli |
| **种属（Species）** | Rattus norvegicus (Rat) |
| **预测分子量（Predicted MW）** | 52.7 kDa |
| **制剂（Form）** | Lyophilized from a 0.2 μm filtered solution of PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and Proclin300. |
| **运输方式（Shipping）** | The product is shipped with ice bag. Upon receipt, store it immediately at the temperature listed below. |
| **稳定性&储存（Stability &Storage）** | 12 months at -20°C to -80°C in lyophilized state.6 months at -20°C to -80°C under sterile conditions after reconstitution.7-10 days at 2°C to 8°C under sterile conditions after reconstitution.Please minimize freeze-thaw cycles. |
| **复溶（Reconstitution）** | Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex. |

## 电泳图（SDS-PAGE image）



Greater than 90% as determined by reducing SDS-PAGE.

## 活性图（Bioactivity image）





**Figure 1. The binding activity of recombinant rat EFNA4 and recombinant human FYN**

Ephrin-A4, also known as EFNA4 and EFL-4, is a member of the ligand of the EPH family. It is mainly expressed in the spleen, lymph nodes, ovary, small intestine and colon of adults, as well as in the heart, lungs, liver, and kidneys of the fetus. It is involved in the development of neurons, blood vessels, and epithelium by regulating cell migration, rejection, and adhesion. Ephrin-A4 has been shown to bind FYN,EphA2, EphA3, EphA4, EphA5, EphA6, EphA7, and EphB1. Thus a functional binding ELISA assay was conducted to detect the interaction of recombinant rat EFNA4 and recombinant human FYN. Briefly, EFNA4 was diluted serially in PBS with 0.01% BSA (pH 7.4). Duplicate samples of 100 µl were then transferred to FYN-coated microtiter wells and incubated for 1h at 37℃. Wells were washed with PBST and incubated for 1h with anti-EFNA4 pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody for 1h at 37℃, wells were aspirated and washed 5 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37℃. Finally, add 50 µL stop solution to the wells and read at 450/630 nm immediately. The binding activity of recombinant rat EFNA4 and recombinant human FYN was shown in Figure 1, the EC50 for this effect is 0.37 ug/mL.

## 注意事项（Note）

For Research Use Only , Not for Diagnostic Use.