
Recombinant Human CD7 (C-6His)

Catalog#:PHH0366

概述（Summary）


别称（Alternative Names）	T-Cell Antigen CD7; GP40; T-Cell Leukemia Antigen; T-Cell Surface Antigen Leu-9; TP41; CD7
纯度（Purity）	Greater than 95% as determined by reducing SDS-PAGE
内毒素（Endotoxin level）	<1 EU/μg as determined by LAL test.
蛋白构建（Construction）	【Ala26-Pro180】-6His
Accession #	P09564
表达宿主（Host）	HEK293
种属（Species）	Human
预测分子量（Predicted MW）	17.47 kDa
制剂（Form）	Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.2.
运输方式（Shipping）	The product is shipped with ice bag. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存（Stability &Storage）	12 months at -20°C to -80°C in lyophilized state.6 months at -20°C to -80°C under sterile conditions after reconstitution.7-10 days at 2°C to 8°C under sterile conditions after reconstitution.Please minimize freeze-thaw cycles.
复溶（Reconstitution）	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图（SDS-PAGE image）



Greater than 95% as determined by reducing SDS-PAGE.

活性图（Bioactivity image）



Immobilized Anti-Human CD7 Antibody (GRI_bio, Research Grade) at 1μg/ml (100 μl/well) can bind Recombinant Human CD7(C-6His).The ED50 of Recombinant Human CD7(C-6His) is 11.39 ng/ml.(Regularly tested).

注意事项（Note）

For Research Use Only , Not for Diagnostic Use.