
Recombinant Human IL-22RA2 (C-6His)

Catalog#:PHH0908

概述（Summary）


别称（Alternative Names）	Interleukin-22 Receptor Subunit Alpha-2; IL-22 Receptor Subunit Alpha-2; IL-22R-Alpha-2; IL-22RA2; Cytokine Receptor Class-II Member 10; Cytokine Receptor Family 2 Member 10; CRF2-10; Cytokine Receptor Family Type 2 Soluble 1; CRF2-S1; Interleukin-22-Binding Protein; IL-22BP; IL22BP; ZcytoR16; IL22RA2
纯度（Purity）	Greater than 95% as determined by reducing SDS-PAGE
内毒素（Endotoxin level）	<1 EU/μg as determined by LAL test.
蛋白构建（Construction）	【Thr22-Pro231】-6His
Accession #	Q969J5-2
表达宿主（Host）	HEK293
种属（Species）	Human
预测分子量（Predicted MW）	25.7 kDa
制剂（Form）	Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.
运输方式（Shipping）	The product is shipped with ice bag. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存（Stability &Storage）	12 months at -20°C to -80°C in lyophilized state.6 months at -20°C to -80°C under sterile conditions after reconstitution.7-10 days at 2°C to 8°C under sterile conditions after reconstitution.Please minimize freeze-thaw cycles.
复溶（Reconstitution）	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图（SDS-PAGE image）



Greater than 95% as determined by reducing SDS-PAGE.

活性图（Bioactivity image）



Immobilized Recombinant Human IL-22 at 1 μg/ml (100 μl/well) can bind Recombinant Human IL-22RA2 (C-6His). The EC50 of Recombinant Human IL-22RA2 (C-6His) is 15.88 ng/ml.(Regularly tested)

注意事项（Note）

For Research Use Only , Not for Diagnostic Use.