
Recombinant Human PD-L1 (C-Flag)

Catalog#:PHH1294

概述（Summary）


别称（Alternative Names）	Programmed Cell Death 1 Ligand 1; PD-L1; PDCD1 Ligand 1; Programmed Death Ligand 1; B7 Homolog 1; B7-H1; CD274; B7H1; PDCD1L1; PDCD1LG1; PDL1
纯度（Purity）	Greater than 95% as determined by reducing SDS-PAGE
内毒素（Endotoxin level）	<1 EU/μg as determined by LAL test.
蛋白构建（Construction）	【Phe19-Thr239】-Flag
Accession #	Q9NZQ7
表达宿主（Host）	HEK293
种属（Species）	Human
预测分子量（Predicted MW）	26.3 kDa
制剂（Form）	Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.
运输方式（Shipping）	The product is shipped with ice bag. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存（Stability &Storage）	12 months at -20°C to -80°C in lyophilized state.6 months at -20°C to -80°C under sterile conditions after reconstitution.7-10 days at 2°C to 8°C under sterile conditions after reconstitution.Please minimize freeze-thaw cycles.
复溶（Reconstitution）	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图（SDS-PAGE image）




Greater than 95% as determined by reducing SDS-PAGE.

活性图（Bioactivity image）



Immobilized Human PD-L1-Flag at 2μg/ml (100 μl/well)can bind Anti-Human PDL1 mAb-Fc.The ED50 of Anti-Human PDL1 mAb-Fc is 14.68 ng/ml.



Loaded Human PD-1-Fc on Protein A Biosensor, can bind Human PD-L1-Flag with an affinity constant of 0.19 uM as determined in BLI assay.

注意事项（Note）

For Research Use Only , Not for Diagnostic Use.