Recombinant Human OX40L (C-mFc)

Catalog#:PHH2044

概述（Summary）

别称（Alternative Names）	Tumor necrosis factor ligand superfamily member 4; Glycoprotein Gp34; OX40 ligand; OX40L; TAX transcriptionally-activated glycoprotein 1; TNFSF4; CD252; TXGP1
纯度（Purity）	Greater than 95% as determined by reducing SDS-PAGE
内毒素（Endotoxin level）	<1 EU/μg as determined by LAL test.
蛋白构建（Construction）	【Gln51-Leu183】-mFc
Accession #	P23510
表达宿主（Host）	HEK293
种属（Species）	Human
预测分子量（Predicted MW）	42.2 kDa
制剂（Form）	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
运输方式（Shipping）	The product is shipped with ice bag. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存（Stability &Storage）	12 months at -20°C to -80°C in lyophilized state.6 months at -20°C to -80°C under sterile conditions after reconstitution.7-10 days at 2°C to 8°C under sterile conditions after reconstitution.Please minimize freeze-thaw cycles.
复溶（Reconstitution）	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

活性图（Bioactivity image）



Immobilized Human OX40-Fc at 8μg/ml (100 μl/well)can bind Human OX40L-mFc.The ED50 of Human OX40L-mFcis 18.54 ng/ml.



Loaded Cynomolgus OX40-His on HIS1K Biosensor, can bind Human OX40L-mFc with an affinity constant of 0.05nM as determined in BLI assay.

注意事项（Note）

For Research Use Only , Not for Diagnostic Use.