
Recombinant Human Siglec-10 (C-Fc)

Catalog#:PHH2149

概述（Summary）


别称（Alternative Names）	SIGLEC10; MGC126774; PRO940; Siglec10; SLG2; sialic acid-binding Ig-like lectin 10; Siglec-10; siglec-like gene 2; Siglec-like protein 2; SLG2sialic acid binding Ig-like lectin 10 Ig-like lectin 7
纯度（Purity）	Greater than 95% as determined by reducing SDS-PAGE
内毒素（Endotoxin level）	<1 EU/μg as determined by LAL test.
蛋白构建（Construction）	【Met17-Thr546】-Fc
Accession #	Q96LC7
表达宿主（Host）	HEK293
种属（Species）	Human
预测分子量（Predicted MW）	85.1 kDa
制剂（Form）	Lyophilized from a 0.2 μm filtered solution of 20mM Tris-HCl, 6% Sucrose, 4% Mannitol, 50mM NaCl, 0.05% Tween 80, pH 8.0.
运输方式（Shipping）	The product is shipped with ice bag. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存（Stability &Storage）	12 months at -20°C to -80°C in lyophilized state.6 months at -20°C to -80°C under sterile conditions after reconstitution.7-10 days at 2°C to 8°C under sterile conditions after reconstitution.Please minimize freeze-thaw cycles.
复溶（Reconstitution）	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图（SDS-PAGE image）



Greater than 95% as determined by reducing SDS-PAGE.

活性图（Bioactivity image）



Immobilized Anti-Human Siglec10 mAb at 2μg/ml (100 μl/well) can bind Human Siglec-10-Fc. The ED50 of Human Siglec-10-Fc is 9.18 ng/ml.

注意事项（Note）

For Research Use Only , Not for Diagnostic Use.