
Recombinant Human IL-17C (C-6His)

Catalog#:PHH2363

概述（Summary）


别称（Alternative Names）	CX2; Cytokine CX2; IL17C; IL-17C; IL-17CMGC126884; interleukin 17C; interleukin-17C
纯度（Purity）	Greater than 95% as determined by reducing SDS-PAGE
内毒素（Endotoxin level）	<1 EU/μg as determined by LAL test.
蛋白构建（Construction）	【His19-Val197】-6His
Accession #	Q9P0M4
表达宿主（Host）	HEK293
种属（Species）	Human
预测分子量（Predicted MW）	20.6 kDa
制剂（Form）	Lyophilized from a 0.2 μm filtered solution of 20mM NaAc, 50mM NaCl, 6% Trehalose, 2% Mannitol, 0.05% Tween 80, pH5.0.
运输方式（Shipping）	The product is shipped with ice bag. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存（Stability &Storage）	12 months at -20°C to -80°C in lyophilized state.6 months at -20°C to -80°C under sterile conditions after reconstitution.7-10 days at 2°C to 8°C under sterile conditions after reconstitution.Please minimize freeze-thaw cycles.
复溶（Reconstitution）	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图（SDS-PAGE image）



Greater than 90% as determined by reducing SDS-PAGE. (QC verified)

活性图（Bioactivity image）



Immobilized Recombinant Human IL-17C (C-6His) at 2μg/ml (100 μl/well) can bind Recombinant Human IL-17RE (C-Fc).The ED50 of Recombinant Human IL-17RE (C-Fc) is 0.60-6.00 ng/ml. (QC verified)

注意事项（Note）

For Research Use Only , Not for Diagnostic Use.