
Active VGF Nerve Growth Factor Inducible (VGF)

Catalog#:PHM12620

概述（Summary）

别称（Alternative Names）	NERP1; NERP2; Neuro-Endocrine Specific Protein VGF; Neuroendocrine Regulatory Peptide
纯度（Purity）	Greater than 95% as determined by reducing SDS-PAGE.
内毒素（Endotoxin level）	Please contact with the lab for this information
蛋白构建（Construction）	【Arg284-Arg576】
Accession #	Q0VGU4
表达宿主（Host）	293F
种属（Species）	Mus musculus (Mouse)
预测分子量（Predicted MW）	34.4 kDa
制剂（Form）	Lyophilized from a 0.2 μm filtered solution of PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and Proclin300.
运输方式（Shipping）	The product is shipped with ice bag. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存（Stability &Storage）	12 months at -20°C to -80°C in lyophilized state.6 months at -20°C to -80°C under sterile conditions after reconstitution.7-10 days at 2°C to 8°C under sterile conditions after reconstitution.Please minimize freeze-thaw cycles.
复溶（Reconstitution）	Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

电泳图（SDS-PAGE image）



Greater than 95% as determined by reducing SDS-PAGE.

活性图（Bioactivity image）




Figure 1. The binding activity of VGF with EGR1

Neurosecretory protein VGF is specifically expressed in a subpopulation of neuroendocrine cells, and is upregulated by nerve growth factor. The protein shares similarities with the secretogranin/chromogranin family, however, its exact function is not known. Besides, Early Growth Response Protein 1 (EGR1) has been identified as an interactor of VGF, thus a binding ELISA assay was conducted to detect the interaction of recombinant mouse VGF and recombinant mouse EGR1. Briefly, EGR1 was diluted serially in PBS with 0.01% BSA (pH 7.4). Duplicate samples of 100µl were then transferred to VGF-coated microtiter wells and incubated for 1h at 37℃. Wells were washed with PBST and incubated for 1h with anti- EGR1 pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody for 1h at 37℃, wells were aspirated and washed 5 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37℃. Finally, add 50µL stop solution to the wells and read at 450/630nm immediately. The binding activity of VGF and EGR1 was shown in Figure 1, the ED50 for this effect is 0.515 ug/mL.

注意事项（Note）

For Research Use Only , Not for Diagnostic Use.