

产品名称: GMP Recombinant Human FGF-10/KGF2
产品货号: PCH90007




概述 (Summary)


英文全称	FGF-10/KGF2
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	≤10 EU/mg
蛋白构建 (Construction)	Recombinant Human FGF-10/KGF2 is produced by our Mammalian cell expression system and the target gene encoding Gln38-Ser208 is expressed.
Accession #	O15520
蛋白标签 (Tag)	Tag free
表达宿主 (Host)	Mammalian cell
种属 (Species)	Human
预测分子量 (Predicted MW)	19.3 kDa
蛋白形态 (Form)	Lyophilized
储存缓冲液 (Buffer)	PBS,5% mannitol and 0.01% Tween 80, pH7.4
运输方式 (Shipping)	The product is shipped with ice bag. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	36 months at -20°C to -80°C in lyophilized state.6 months at -20°C to -80°C under sterile conditions after reconstitution.7-10 days at 2°C to 8°C under sterile conditions after reconstitution.Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

产品名称: GMP Recombinant Human FGF-10/KGF2
产品货号: PCH90007



活性验证图 (Bioactivity image)



Measured by its ability to induce IL-11 secretion by Saos-2 human osteosarcoma cells. The ED50 for this effect is $\leq 2 \mu\text{g/mL}$.

背景 (Background)

分子别名 (Alternative Names)

Fibroblast growth factor 10; FGF-10; Keratinocyte growth factor 2; FGF10; KGF-2; KGF2

背景介绍 (References)

Fibroblast growth factor 10 (FGF-10, KGF-2), is a member of the fibroblast growth factor (FGF) family that includes FGF-3, -7, and -22. KGF-2 is secreted by mesenchymal cells and associates with extracellular FGF-BP. It preferentially binds and activates epithelial cell FGFR2 and interacts more weakly with FGFR1. It plays an important role in the regulation of embryonic development, cell proliferation and cell differentiation. It exhibits mitogenic activity for keratinizing epidermal cells, but essentially no activity for fibroblasts, which is similar to the biological activity of FGF7. FGF10 is required for normal branching morphogenesis. Defects in FGF10 are the

产品名称: GMP Recombinant Human FGF-10/KGF2
产品货号: PCH90007



cause of autosomal dominant aplasia of lacrimal and salivary glands (ALSG). ALSG has variable expressivity, and affected individuals may have aplasia or hypoplasia of the lacrimal, parotid, submandibular and sublingual glands and absence of the lacrimal puncta. The disorder is characterized by irritable eyes, recurrent eye infections, epiphora (constant tearing) and xerostomia (dryness of the mouth), which increases the risk of dental erosion, dental caries, periodontal disease and oral infections.

注意事项 (Note)

For research use only .