


产品名称: GMP Recombinant Human OSM
产品货号: PCH90036



概述 (Summary)

英文全称	OSM
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	≤ 10 EU/mg
蛋白构建 (Construction)	Recombinant Human OSM is produced by our Mammalian cell expression system and the target gene encoding Ala26-Arg221 is expressed.
Accession #	P13725
蛋白标签 (Tag)	Tag free
表达宿主 (Host)	Mammalian cell
种属 (Species)	Human
预测分子量 (Predicted MW)	22 kDa
蛋白形态 (Form)	Lyophilized
储存缓冲液 (Buffer)	PBS, 5% mannitol and 0.01% Tween 80, pH7.4
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	36 months at -20°C to -80°C in lyophilized state. 6 months at -20°C to -80°C under sterile conditions after reconstitution. 7-10 days at 2°C to 8°C under sterile conditions after reconstitution. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than $100\mu\text{g/ml}$. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.


电泳图 (SDS-PAGE image)



产品名称: GMP Recombinant Human OSM
产品货号: PCH90036



活性验证图 (Bioactivity image)



The ED50 for this effect is ≤ 2 ng/mL.

背景 (Background)

分子别名 (Alternative Names)

Oncostatin-M; OSM

背景介绍 (References)

Oncostatin M (OSM) is a glycoprotein belonging to the interleukin-6 family of cytokines that includes leukemia-inhibitory factor, granulocyte colony-stimulating factor, and interleukin 6. OSM encodes a growth regulator, which inhibits the proliferation of a number of tumor cell lines. It stimulates proliferation of AIDS-KS cells. OSM regulates cytokine production, including IL-6, G-CSF and GM-CSF from endothelial cells. OSM is considered as a pleiotropic cytokine that initiates its biological activities through specific cell surface receptors. The low affinity LIF receptor that shares the similarity of containing protein gp130 has now been identified to be a component of a high-affinity OSM receptor that will transduce OSM signals. OSM has also been shown to play a role in both pro and anti-inflammatory actions. OSM may also be involved in many biometabolism processes including liver development, haematopoiesis, inflammation, bone formation and destruction and possibly CNS development.

注意事项 (Note)

For research use only .