


概述 (Summary)

英文全称	Annexin A2/ANXA2
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Human Annexin A2 is produced by our E.coli expression system and the target gene encoding Ser2-Asp339 is expressed.
Accession #	P07355
蛋白标签 (Tag)	
表达宿主 (Host)	E.coli
种属 (Species)	Human
预测分子量 (Predicted MW)	38.6 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of 20mM Histidine-HCl, 6% Sucrose, 4% Mannitol, 0.05%Tween 80, pH 5.5.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Human ANXA2
Catalog #: PEH0071



背景 (Background)

分子别名 (Alternative Names)

Annexin A2; Annexin II; Annexin-2; Calpactin I Heavy Chain; Calpactin-1 Heavy Chain; Chromobindin-8; Lipocortin II; Placental Anticoagulant Protein IV; PAP-IV; Protein I; p36; ANXA2; ANX2; ANX2L4; CAL1H; LPC2D

背景介绍 (References)

Annexin A2 (ANXA2) is a member of the annexin family and has roles in the regulation of cellular growth and in signal transduction pathways. ANXA2 protein is associated with sickle cell osteonecrosis and the expression reduce of ANXA2 is associated with osteosarcoma metastases. ANXA2 functions as an autocrine factor, it can increases osteoclast formation and bone resorption. ANXA2 is involved in muscular dystrophies. In humans, the up-regulation of ANXA2 is related with colon adenocarcinoma cell differentiation.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.