


概述 (Summary)

英文全称	ATP-binding cassette sub-family B member 5/ABCB5
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Human ATP-binding Cassette Sub-family B Member 5 is produced by our E.coli expression system and the target gene encoding Ile141-Val247 is expressed with a Trx tag at the N-terminus.
Accession #	Q2M3G0
蛋白标签 (Tag)	
表达宿主 (Host)	E.coli
种属 (Species)	Human
预测分子量 (Predicted MW)	29.4 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Human ABCB5 (N-Trx)
Catalog #: PEH0113



背景 (Background)

分子别名 (Alternative Names)

ATP-binding cassette sub-family B member 5; P-glycoprotein ABCB5; ABCB5 P-gp; ABCB5;

背景介绍 (References)

ATP-binding cassette sub-family B member 5(ABCB5) is a plasma membrane-spanning protein. ABCB5 is principally expressed in physiological skin and human malignant melanoma. ABCB5 has been suggested to regulate skin progenitor cell fusion and mediate chemotherapeutic drug resistance in stem-like tumor cell subpopulations in human malignant melanoma. It is commonly over-expressed on circulating melanoma tumour cells. Furthermore, the ABCB5+ melanoma- initiating cells were demonstrated to express FLT1 (VEGFR1) receptor tyrosine kinase which was functionally required for efficient xenograft tumor formation, as demonstrated by shRNA knockdown experiments.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.