


Product Name: Recombinant Human B3GAT3 (C-6His)
Catalog #: PEH0117



概述 (Summary)

英文全称	B3GAT3/GLCATI
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Human Galactosylgalactosylxylosylprotein 3-beta-glucuronosyltransferase 3 is produced by our E.coli expression system and the target gene encoding Glu72-Val335 is expressed with a 6His tag at the C-terminus.
Accession #	O94766
蛋白标签 (Tag)	
表达宿主 (Host)	E.coli
种属 (Species)	Human
预测分子量 (Predicted MW)	30.4 KDa
蛋白形态 (Form)	Supplied as a 0.2 μm filtered solution of 20mM Tris-HCl, 150mM NaCl, 2mM EDTA, 20% Glycerol, pH 8.0.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped on dry ice/polar packs. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	

电泳图 (SDS-PAGE image)



背景 (Background)

Product Name: Recombinant Human B3GAT3 (C-6His)
Catalog #: PEH0117



分子别名 (Alternative Names)

Unconjugated

背景介绍 (References)

Galactosylgalactosylxylosylprotein 3-beta-glucuronosyltransferase 3 (B3GAT3) is an enzyme that in humans is encoded by the B3GAT3 gene, belongs to the glycosyltransferase 43 family. B3GAT3 is involved in a number of biological processes such as catalyzing the formation of the glycosaminoglycan-protein linkage by way of a glucuronyl transfer reaction in the final step of the biosynthesis of the linkage region of proteoglycans, forming the linkage tetrasaccharide present in heparan sulfate and chondroitin sulfate, gGlycosaminoglycans biosynthesis, transferring a glucuronic acid moiety from the uridine diphosphate-glucuronic acid (UDP-GlcUA) to the common linkage region trisaccharide Gal-beta-1,3-Gal-beta-1,4-Xyl covalently bound to a Ser residue at the glycosaminylglycan attachment site of proteoglycans. It also plays a role in the biosynthesis of I2/HNK-1 carbohydrate epitope on glycoproteins, shows strict specificity for Gal-beta-1,3-Gal-beta-1,4-Xyl, exhibiting negligible incorporation into other galactoside substrates including Galbeta1-3Gal beta1-O-benzyl, Galbeta1-4GlcNAc and Galbeta1-4Glc and stimulates 2-phosphoxylose phosphatase activity of PXYLP1 in presence of uridine diphosphate-glucuronic acid (UDP-GlcUA) during completion of linkage region formation.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.