

Product Name: Recombinant Human CRYAA (C-6His)
Catalog #: PEH0453




概述 (Summary)

英文全称	CRYAA/Alpha-crystallin A chain
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Human Alpha-Crystallin A Chain is produced by our E.coli expression system and the target gene encoding Met1-Ser173 is expressed with a 6His tag at the C-terminus.
Accession #	P02489
蛋白标签 (Tag)	
表达宿主 (Host)	E.coli
种属 (Species)	Human
预测分子量 (Predicted MW)	20.9 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of PBS, 2mM EDTA, pH 8.0.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Human CRYAA (C-6His)
Catalog #: PEH0453



背景 (Background)

分子别名 (Alternative Names)

Alpha-Crystallin A Chain; Heat Shock Protein Beta-4; HspB4; Alpha-Crystallin A Chain; Short Form; CRYAA; CRYA1; HSPB4

背景介绍 (References)

Alpha-Crystallin A Chain (CRYAA) belongs to the small heat shock protein (HSP20) family and can be induced by heat shock. The expression of CRYAA is preferentially restricted to the lens cell. CRYAA may contribute to the transparency and refractive index of the lens. CRYAA has chaperone-like activity, preventing aggregation of various proteins under a wide range of stress conditions. Two additional functions of CRYAA are an autokinase activity and participation in the intracellular architecture.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.