Product Name: Recombinant Human EDF1 (C-6His)

Catalog #: PEH0554



概述 (Summary)

英文全称 EDF1/MBF1

纯度 (Purity) Greater than 95% as determined by reducing SDS-PAGE

内毒素 (Endotoxin level) <1 EU/μg as determined by LAL test.

蛋白构建 (Construction) Recombinant Human Endothelial Differentiation-Related Factor 1 is

produced by our E.coli expression system and the target gene encoding

Ala2-Lys148 is expressed with a 6His tag at the C-terminus.

Accession # O60869

蛋白标签 (Tag)

表达宿主 (Host)E.coli种属 (Species)Human预测分子量 (Predicted MW)17.4 KDa

蛋白形态 (Form) Lyophilized from a 0.2 μm filtered solution of 20mM Tris-HCl, 150mM NaCl,

pH 8.0.

储存缓冲液 (Buffer)

运输方式 (Shipping) The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

稳定性&储存 (Stability &Storage) Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3

months under sterile conditions after opening. Please minimize freeze-thaw

cycles.

复溶 (Reconstitution) Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It

is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized

protein in distilled water. Please aliquot the reconstituted solution to minimize

freeze-thaw cycles.


电泳图 (SDS-PAGE image)

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838

Product Name: Recombinant Human EDF1 (C-6His)

Catalog #: PEH0554





背景 (Background)

分子别名 (Alternative Names)

背景介绍 (References)

Endothelial Differentiation-Related Factor 1; EDF-1; Multiprotein-Bridging Factor 1; MBF1; EDF1

Endothelial Differentiation-Related Factor 1 (EDF1) is a 148 amino acid transcriptional coactivator that contains 1 HTH cro/C1-type DNA-binding domain. It has been postulated that the protein functions as a bridging molecule that interconnects regulatory proteins and the basal transcriptional machinery, thereby modulating the transcription of genes involved in endothelial differentiation. When endothelial cells are induced to differentiate in vitro, EDF1 is downregulated, leading to inhibition of cell growth and cell polarization. EDF1 binds calmodulin thorough its IQ domain and regulates nitric oxide synthase activity through calmodulin sequestration in the cytoplasm. Though ubiquitously expressed, EDF1 is most abundant in adult liver, heart, adipose tissues, intestine and pancreas. In fetal tissues, EDF1 is most abundant in kidney. There are two isoforms of EDF1 that are produced as a result of alternative splicing events.

注意事项 (Note)

For Research Use Only, Not for Diagnostic Use.