


概述 (Summary)

英文全称	HDHD2
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Human Haloacid Dehalogenase-Like Hydrolase Domain-Containing Protein 2 is produced by our E.coli expression system and the target gene encoding Met1-Leu259 is expressed with a 6His tag at the N-terminus.
Accession #	Q9H0R4
蛋白标签 (Tag)	
表达宿主 (Host)	E.coli
种属 (Species)	Human
预测分子量 (Predicted MW)	30.7 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of 20mM Tris-HCl, 50mM NaCl, pH 8.0.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Human HDHD2 (N-6His)
Catalog #: PEH0777



背景 (Background)

分子别名 (Alternative Names)

Haloacid Dehalogenase-Like Hydrolase Domain-Containing Protein 2; HDHD2

背景介绍 (References)

Haloacid Dehalogenase-Like Hydrolase Domain-Containing Protein 22 (HDHD2) is a member of the HAD-like hydrolase superfamily. HDHD2 includes L-2-Haloacid Dehalogenase, Epoxide Hydrolases and Phosphatases. There are two active sites in HDHD2 - an L-2-Haloacid Dehalogenase and a Carboxylate group. The L-2-Haloacid Dehalogenase active site catalyzes the hydrolytic dehalogenation of D- and L-2-Haloalkanoic Acids, producing L- and D-2-Hydroxyalkanoic Acids.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.