


概述 (Summary)

英文全称	Krueppel-like factor 6/KLF6
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Human Krueppel-like Factor 6 is produced by our E.coli expression system and the target gene encoding Met1-Ser109 is expressed.
Accession #	Q99612-3
蛋白标签 (Tag)	
表达宿主 (Host)	E.coli
种属 (Species)	Human
预测分子量 (Predicted MW)	12.6 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Human KLF6
Catalog #: PEH1052



背景 (Background)

分子别名 (Alternative Names)

Krueppel-Like Factor 6; B-Cell-Derived Protein 1; Core Promoter Element-Binding Protein; GC-Rich Sites-Binding Factor GBF; Proto-Oncogene BCD1; Suppressor of Tumorigenicity 12 Protein; Transcription Factor Zf9; KLF6; BCD1; COPEB; CPBP; ST12

背景介绍 (References)

Krueppel-Like Factor 6 (KLF6) belongs to the krueppel C2H2-type zinc-finger protein family. KLF6 contains three C2H2-type zinc fingers and localizes in the nucleus. KLF6 expression is highest in the placenta followed by spleen, thymus, prostate, testis, small intestine and colon. However, it is weakly expressed in the pancreas, lung, liver, heart, and skeletal muscle. KLF6 functions as a transcriptional activator and could play a role in B-cell growth and development. Defects in KLF6 will result in gastric cancer and prostate cancer.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.