


Product Name: Recombinant Human MDH1 (C-6His)
Catalog #: PEH1142



概述 (Summary)

英文全称	MDH1/Malate dehydrogenase, cytoplasmic
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Human Malate Dehydrogenase, Cytoplasmic is produced by our E.coli expression system and the target gene encoding Ser2-Ala334 is expressed with a 6His tag at the C-terminus.
Accession #	P40925
蛋白标签 (Tag)	
表达宿主 (Host)	E.coli
种属 (Species)	Human
预测分子量 (Predicted MW)	37.5 KDa
蛋白形态 (Form)	Supplied as a 0.2 μm filtered solution of 20mM Tris-HCl, 150mM NaCl, pH 8.0.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped on dry ice/polar packs. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	

电泳图 (SDS-PAGE image)



背景 (Background)

Product Name: Recombinant Human MDH1 (C-6His)
Catalog #: PEH1142



分子别名 (Alternative Names)

Malate Dehydrogenase Cytoplasmic; Cytosolic Malate Dehydrogenase; Diiodophenylpyruvate Reductase; MDH1; MDHA

背景介绍 (References)

Malate Dehydrogenase, Cytoplasmic (MDH1) is an enzyme which belongs to the MDH Type 2 sub-family of LDH/MDH superfamily. MDH1 is involved in the Citric Acid Cycle that catalyzes the conversion of Malate into Oxaloacetate (using NAD⁺) and vice versa. MDH1 should not be confused with Malic Enzyme, which catalyzes the conversion of Malate to Pyruvate, producing NADPH. MDH1 also participates in Gluconeogenesis, the synthesis of Glucose from smaller molecules. Pyruvate in the mitochondria is acted upon by Pyruvate Carboxylase to form Oxaloacetate, a Citric Acid Cycle intermediate. In order to transport the Oxaloacetate out of the Mitochondria, Malate Dehydrogenase reduces it to Malate, and it then traverses the inner Mitochondrial membrane. Once in the cytosol, the Malate is oxidized back to Oxaloacetate by MDH1. Finally, Phosphoenol-Pyruvate Carboxy Kinase (PEPCK) converts Oxaloacetate to Phosphoenol Pyruvate.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.