

产品名称: Recombinant Human NAMPT (N-6His)
产品货号: PEH1362




概述 (Summary)

英文全称	Pre-B-Cell Colony-Enhancing Factor 1/PBEF/Visfatin/NAMPT/Nicotinamide phosphoribosyltransferase
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Human Pre-B-Cell Colony-Enhancing Factor 1 is produced by our E.coli expression system and the target gene encoding Met1-His491 is expressed with a 6His tag at the N-terminus.
Accession #	P43490
表达宿主 (Host)	E.coli
种属 (Species)	Human
预测分子量 (Predicted MW)	57 KDa
制剂 (Form)	Lyophilized from a 0.2 μm filtered solution of 20mM PB, 10% Trehalose, 100mM Arg-HCl, 0.02% Tween 80, pH6.0.
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

产品名称: Recombinant Human NAMPT (N-6His)
产品货号: PEH1362



背景 (Background)

分子别名 (Alternative Names)

Pre-B cell-enhancing factor; Nicotinamide phosphoribosyltransferase; NAmPRTase; Nampt; Pre-B-cell colony-enhancing factor 1; Visfatin; NAMPT; PBEF; PBEF1

背景介绍 (References)

Pre-B cell colony enhancing factor (PBEF) was originally identified as a cytokine that potentiated the clonal expansion and differentiation of pre-B cells, but it is also acknowledged to be the ubiquitous intracellular enzyme nicotinamide phosphoribosyltransferase (NAMPT) and the adipokine "visfatin". PBEF is constitutively expressed in the fetal membranes where its greatest expression is in the amnion. It has intracellular and extracellular forms. Most of the intracellular functions of PBEF are due to its role as a Nampt which can induce angiogenesis through upregulation of VEGF and VEGFR and secretion of MCP-1. Extracellular PBEF has been shown to increase inflammatory cytokines, such as TNF- α , IL-1 β , IL-16, and TGF- β 1. PBEF also increases the production of IL-6, TNF- α , and IL-1 β in CD14⁺ monocytes, macrophages, and dendritic cells, enhances the effectiveness of T cells.

注意事项 (Note)

For research use only .