

产品名称: Recombinant Human PFN2
产品货号: PEH1370




概述 (Summary)

英文全称	Profilin-2/PFN2
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/ μ g as determined by LAL test.
蛋白构建 (Construction)	Recombinant Human Profilin-2 is produced by our E.coli expression system and the target gene encoding Met1-Phe140 is expressed.
Accession #	P35080
表达宿主 (Host)	E.coli
种属 (Species)	Human
预测分子量 (Predicted MW)	15 KDa
制剂 (Form)	Lyophilized from a 0.2 μ m filtered solution of 20mM Tris-HCl, 150mM NaCl, pH 8.0.
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Lyophilized protein should be stored at $\leq -20^{\circ}\text{C}$, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8 $^{\circ}\text{C}$ for 2-7 days. Aliquots of reconstituted samples are stable at $\leq -20^{\circ}\text{C}$ for 3 months.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 μ g/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 μ g/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

产品名称: Recombinant Human PFN2
产品货号: PEH1370



背景 (Background)

分子别名 (Alternative Names)

Profilin-II; PFN2; Profilin-2; PFL

背景介绍 (References)

Profilin-II (PFN2) is ubiquitous protein which belongs to the profilin family. PFN2 binds to actin, then affects the structure of the cytoskeleton. At high concentrations, profilin prevents the polymerization of actin, while increases that at low concentrations. PFN2 is a ubiquitous actin monomer-binding protein. It regulates actin polymerization in response to extra cellular signals. PFN2 binds to PIP2; it inhibits the formation of IP3 and DG.

注意事项 (Note)

For research use only .