


概述 (Summary)

英文全称	SNCA/Alpha-synuclein
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<0.001 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Human Alpha-Synuclein is produced by our E.coli expression system and the target gene encoding Met1-Ala140 is expressed.
Accession #	P37840
蛋白标签 (Tag)	
表达宿主 (Host)	E.coli
种属 (Species)	Human
预测分子量 (Predicted MW)	14.46 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of 20 mM Tris-HCl, 15% Maltose, 0.1% Tween 80, 10 mM Methionine, pH 8.0.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Human SNCA
Catalog #: PEH1544



背景 (Background)

分子别名 (Alternative Names)

Alpha-Synuclein; Non-A Beta Component of AD Amyloid; Non-A4 Component of Amyloid Precursor; NACP; SNCA; NACP; PARK1

背景介绍 (References)

Alpha-Synuclein (SNCA) is a member of the Synuclein family. SNCA is expressed principally in brain but also expressed in low concentrations in all tissues except liver. SNCA interacts with UCHL1, Phospholipase D and histones. SNCA can include beta- and gamma-synuclein. In addition, SNCA is an important regulatory component of vesicular transport in neuronal cells. It has been suggested that SNCA is related to the pathogenesis of Parkinsons Disease and neurodegenerative disorders. Defects in SNCA will lead to Dementia Lewy Body (DLB).

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.