


概述 (Summary)

英文全称	STAT6
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/ μ g as determined by LAL test.
蛋白构建 (Construction)	Recombinant Human Signal Transducer And Activator Of Transcription 6 is produced by our E.coli expression system and the target gene encoding Ser627-Ser837 is expressed with a 6His tag at the C-terminus.
Accession #	P42226
蛋白标签 (Tag)	
表达宿主 (Host)	E.coli
种属 (Species)	Human
预测分子量 (Predicted MW)	23.9 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μ m filtered solution of PBS, pH 7.4.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at $\leq -70^{\circ}\text{C}$, stable for 6 months after receipt. Store at $\leq -70^{\circ}\text{C}$, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 μ g/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 μ g/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Human STAT6 (C-6His)
Catalog #: PEH1570



背景 (Background)

分子别名 (Alternative Names)

Signal Transducer and Activator of Transcription 6; IL-4 Stat; STAT6

背景介绍 (References)

Signal Transducer and Activator of Transcription 6 (STAT6) is a member of the STAT family of transcription factors. At least seven STATs exist: STAT1, 2, 3, 4, 5a, 5b, and 6. They are responsible for an array of cellular activities including regulating growth, survival, differentiation, motility, and the immune response. STAT6 plays a central role in exerting IL4 mediated biological responses. It is found to induce the expression of BCL2L1/BCL-X(L), which is responsible for the anti-apoptotic activity of IL4. Knockout studies in mice suggested the roles of this gene in differentiation of T helper 2 (Th2) cells, expression of cell surface markers, and class switch of immunoglobulins. STAT6 has been shown to interact with EP300, CREB-binding protein, NFKB1, Nuclear receptor coactivator 1, IRF4 and SND1.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.