


概述 (Summary)

英文全称	Tachykinin-3/TAC3
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Human Tachykinin-3 is produced by our E.coli expression system and the target gene encoding Gln17-Glu121 is expressed with a 6His tag at the N-terminus.
Accession #	Q9UHF0
蛋白标签 (Tag)	
表达宿主 (Host)	E.coli
种属 (Species)	Human
预测分子量 (Predicted MW)	13.9 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of 20mM Tris-HCl, pH 8.0.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Human TAC3 (N-6His)
Catalog #: PEH1603



背景 (Background)

分子别名 (Alternative Names)

Tachykinin-3; ZNEUROK1; Neurokinin-B; NKB; Neuromedin-K; TAC3; NKNB; UNQ585/PRO1155

背景介绍 (References)

Tachykinin 3 (TAC3) is a secreted protein that belongs to the Tachykinin family. Tachykinins are active peptides that excite neurons and evoke behavioral responses; they are potent vasodilators and secretagogues, and contract many smooth muscles in pregnancy. TAC3 is primarily expressed in the central and peripheral nervous systems and functions as a neurotransmitter. It is also expressed in the outer syncytiotrophoblast of the placenta and may be associated with pregnancy-induced hypertension and pre-eclampsia. TAC3 acts as the ligand for the neurokinin-3 receptor, mutations in this gene are associated with normosmic hypogonadotropic hypogonadism.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.