


概述 (Summary)

英文全称	Thioredoxin/TXN/Trx
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Human Thioredoxin is produced by our E.coli expression system and the target gene encoding Met1-Val105 is expressed with a 6His tag at the N-terminus.
Accession #	P10599
蛋白标签 (Tag)	
表达宿主 (Host)	E.coli
种属 (Species)	Human
预测分子量 (Predicted MW)	13.9 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of 20mM PB, 1mM EDTA, 2mM DTT, pH 7.2.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Human TXN (N-6His)
Catalog #: PEH1632



背景 (Background)

分子别名 (Alternative Names)

Thioredoxin; Trx; ATL-Derived Factor; ADF; Surface-Associated Sulphydryl Protein; SASP; TXN; TRDX; TRX; TRX1

背景介绍 (References)

Thioredoxin (TXN) is a member of the Thioredoxin family. Thioredoxin exists as a disulfide-linked homodimer and contains one Thioredoxin domain. Thioredoxin is up-regulated by ionizing radiation. Thioredoxin participates in various redox reactions through the reversible oxidation of its active center dithiol to a disulfide and catalyzes dithiol-disulfide exchange reactions. Thioredoxin also plays a role in the reversible S-nitrosylation of cysteine residues in target proteins, and thereby contributes to the response to intracellular nitric oxide.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.