

Product Name: Recombinant Human TXLNA (N, C-6His)
Catalog #: PEH1750




概述 (Summary)

英文全称	TXLNA/ α -taxilin
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/ μ g as determined by LAL test.
蛋白构建 (Construction)	Recombinant Human Alpha-Taxilin is produced by our E.coli expression system and the target gene encoding Met1-Lys162 is expressed with a 6His tag at the N-terminus, 6His tag at the C-terminus.
Accession #	P40222
蛋白标签 (Tag)	
表达宿主 (Host)	E.coli
种属 (Species)	Human
预测分子量 (Predicted MW)	20.4 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μ m filtered solution of PBS, pH 7.4.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at $\leq -70^{\circ}\text{C}$, stable for 6 months after receipt. Store at $\leq -70^{\circ}\text{C}$, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 μ g/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 μ g/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Human TXLNA (N, C-6His)
Catalog #: PEH1750



背景 (Background)

分子别名 (Alternative Names)

Alpha-Taxilin; TXLNA; TXLN

背景介绍 (References)

α -Taxilin belongs to the taxilin family. α -Taxilin exists in almost all tissues, with higher expression levels observed in the heart, kidney, liver, and pancreas. α -Taxilin binds to the C-terminal coiled coil region of syntaxin family members STX1A, STX3A, and STX4A, but not when these proteins are complexed with SNAP25, VAMP2 or STXBP1, suggesting that it interacts with syntaxins that do not form the SNARE complex. It is shown that α -Taxilin plays multiple roles in the generation and maintenance of neurons through modulation of the NAC-mediated translational machinery and/or the syntaxin-mediated vesicle traffic in the soma. In addition, α -Taxilin may be involved in intracellular vesicle traffic and potentially in calcium-dependent exocytosis in neuroendocrine cells.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.