


概述 (Summary)

英文全称	Zinc finger protein 225/EGR1/ZNF225
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Human Early Growth Response Protein 1 is produced by our E.coli expression system and the target gene encoding Gln282-Ser433 is expressed with a 6His tag at the N-terminus.
Accession #	P18146
蛋白标签 (Tag)	
表达宿主 (Host)	E.coli
种属 (Species)	Human
预测分子量 (Predicted MW)	19.9 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Human EGR1 (N-6His)
Catalog #: PEH1843



背景 (Background)

分子别名 (Alternative Names)

EGR-1; Early growth response protein 1; Zif268; zinc finger protein 225; NGFI-A ; nerve growth factor-induced protein A;

背景介绍 (References)

EGR-1 belongs to the EGR family of C2H2-type zinc finger proteins. It is a nuclear protein and functions as a transcriptional regulator. EGR-1 recognizes and binds to the DNA sequence 5-CGCCCCCGC-3(EGR-site).The products of target genes it activates are required for differentiation and mitogenesis. Studies suggest this is a tumor suppressor gene. EGR-1 has a distinct pattern of expression in the brain, and its induction has been shown to be associated with neuronal activity. Several studies suggest it has a role in neuronal plasticity. EGR-1 has also been found to regulate the expression of synaptobrevin II (a protein important for synaptic exocytosis).

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.