


概述 (Summary)

英文全称	SAA2/Serum Amyloid A2
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Human Serum Amyloid A-2 Protein is produced by our E.coli expression system and the target gene encoding Arg19-Tyr122 is expressed with a 6His tag at the N-terminus.
Accession #	AAH20795.1
蛋白标签 (Tag)	
表达宿主 (Host)	E.coli
种属 (Species)	Human
预测分子量 (Predicted MW)	13.2 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, 1mM EDTA, pH 7.4.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Human SAA2 (N-6His)
Catalog #: PEH1962



背景 (Background)

分子别名 (Alternative Names)

SAA; SAA2; Serum Amyloid A2

背景介绍 (References)

Serum amyloid A-2 protein (SAA2) belongs to the SAA family. It expressed by the liver and secreted in plasma. SAA2 functions as major acute phase reactant and could works as apolipoprotein of the HDL complex. Increased levels of A-SAA in serum are indicative of inflammatory disease. When highly expressed, SAA can displace ApoA1 as the major apolipoprotein in HDL complexes, weakening the function of HDL as a reverse (lipid clearing) cholesterol transporter. A highly charged region of SAA2 and SAA1 (aa 36-68) contains putative fibronectin- and laminin-binding motifs. This region also binds heparin sulfate proteoglycans at mildly acidic pH and promotes aggregation of A-SAA.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.