


概述 (Summary)

英文全称	Betacellulin/Btc
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Mouse Betacellulin is produced by our E.coli expression system and the target gene encoding Asp32-Gln118 is expressed with a 6His tag at the N-terminus.
Accession #	Q543J8
蛋白标签 (Tag)	
表达宿主 (Host)	E.coli
种属 (Species)	Mouse
预测分子量 (Predicted MW)	12.2 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of 20mM Citrate, 10% Trehalose, 2% Mannitol, 0.05% Tween80, pH 4.5.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)



背景 (Background)

分子別名 (Alternative Names)

Btc;Betacellulin;Betacellulin; epidermal growth factor family member;Betacellulin; epidermal growth factor family member; isoform CRA_a;mCG_12529

背景介绍 (References)

Mouse Betacellulin is a single type I membrane protein which belongs to the EGF family of cytokines. EGF family has many members including EGF, TGF- α , Amphiregulin, HB-EGF, Epiregulin, Tomoregulin and the Neuregulins. Betacellulin is characterised by a six-cysteine consensus motif that forms three intra-molecular disulfide bonds crucial for binding the ErbB receptor family. Betacellulin is expressed in several tissues and tumor cells including kidney, uterus, liver, pancreas and small intestine. Betacellulin binds and activates ErbB-1 and ErbB-4 homodimers. Betacellulin is thought to play a role in the differentiation of pancreatic beta cells. Human and mouse mature BTC protein are 80% identical at the amino acid sequence level. Betacellulin is involved in many biological processes such as stimulating gastrointestinal growth. It is proteolytically processed from a larger membrane-anchored precursor and is a potent mitogen for a wide variety of cell types.

注意事项 (Note)

For Research Use Only, Not for Diagnostic Use.