


概述 (Summary)

英文全称	FGF-9/Fibroblast Growth Factor 9
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<0.01 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Mouse Fibroblast Growth Factor 9 is produced by our E.coli expression system and the target gene encoding Met1/xadSer208 is expressed with a 6His tag at the N-terminus.
Accession #	P54130
蛋白标签 (Tag)	
表达宿主 (Host)	E.coli
种属 (Species)	Mouse
预测分子量 (Predicted MW)	24.4 KDa
蛋白形态 (Form)	Supplied as a 0.2 μm filtered solution of 20mM Tris-HCl, 150mM NaCl, 5% Trehalose, 1mM EDTA, 20% Glycerol, 1mM DTT, pH 8.5.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped on dry ice/polar packs. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	0.00.0

电泳图 (SDS-PAGE image)



背景 (Background)

分子别名 (Alternative Names) Fibroblast growth factor 9; FGF-9; Glia-activating factor; GAF; heparin-binding growth factor-9; HBGF-9; Fgf9; Fgf-9

背景介绍 (References) Fibroblast growth factor-9 (FGF-9) is an approximately 26 kDa secreted glycoprotein of the FGF family. Secreted mouse FGF-9 lacks the N-terminal 1-3 aa and shares //u003E98% sequence identity with rat, human, equine, porcine and bovine FGF-9. FGF-9 plays an important role in the regulation of embryonic development, cell proliferation, cell differentiation and cell migration. In the mouse embryo the location and timing of FGF-9 expression affects development of the skeleton, cerebellum, lungs, heart, vasculature, digestive tract, and testes .It may have a role in glial cell growth and differentiation during development, gliosis during repair and regeneration of brain tissue after damage, differentiation and survival of neuronal cells, and growth stimulation of glial tumors. Deletion of mouse FGF-9 is lethal at birth due to lung hypoplasia, and causes rhizomelia, or shortening of the proximal skeleton. An unusual constitutive dimerization of FGF 9 buries receptor interaction sites which lowers its activity, and increases heparin affinity which inhibits diffusion. A spontaneous mouse mutant, Eks, interferes with dimerization, resulting monomeric, diffusible FGF-9 that causes elbow and knee synostoses (joint fusions) due to FGF-9 misexpression in developing joints.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.