


概述 (Summary)

英文全称	IL-33/Interleukin-33/IL-1F11/Interleukin-1 Family Member 11/Nuclear Factor From High Endothelial Venules/NF-HEV/C9orf26/NFHEV
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<0.01 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Mouse Interleukin-33 is produced by our E.coli expression system and the target gene encoding Ser109-Ile266 is expressed.
Accession #	Q8BVZ5
蛋白标签 (Tag)	
表达宿主 (Host)	E.coli
种属 (Species)	Mouse
预测分子量 (Predicted MW)	17.6 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of PBS, 1mM DTT, pH 7.4.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Mouse IL-33
Catalog #: PEM0924



背景 (Background)

分子别名 (Alternative Names)

Interleukin 33; IL-33; IL33; C9orf26; NKHEV; Interleukin-1 family member 11; DVS27; NF-HEV and IL- 1F11

背景介绍 (References)

Mouse Interleukin 33 (IL-33) is a 30 kDa proinflammatory cytokine which may also regulates gene transcription in producer cells. IL-33 is constitutively expressed in smooth muscle and airway epithelia. IL-33 was identified based on sequence and structural homology with IL-1 family cytokines. It is up-regulated in arterial smooth muscle, dermal fibroblasts, and keratinocytes following IL-1 alpha or IL-1 beta stimulation. IL-33 is structurally related to IL-1, which induces helper T cells to produce type 2 cytokines and acts through the receptor IL1RL-1. Binding IL-33 to this receptor activates NF-kappa-B and MAP kinases and induces in vitro Th2 cells to produce cytokines. In vivo, IL-33 induces the expression of IL-4, IL-5, IL-13 and also leads to severe pathological changes in mucosal organs.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.