


概述 (Summary)

英文全称	IL-36 beta/IL-36b/IL1F8
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Mouse Interleukin-36 Beta is produced by our E.coli expression system and the target gene encoding Ser31-Lys183 is expressed.
Accession #	Q9D6Z6
蛋白标签 (Tag)	
表达宿主 (Host)	E.coli
种属 (Species)	Mouse
预测分子量 (Predicted MW)	17.6 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of 20mM Tris-HCl, 150mM NaCl, 1mM EDTA, pH 8.0.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Mouse IL-36b
Catalog #: PEM0926



背景 (Background)

分子别名 (Alternative Names)

Il36b;Interleukin-36 beta;Interleukin-1 family member 8;IL-1F8;Fil1e; Il1f8

背景介绍 (References)

Mouse Interleukin 36 beta (IL-36B) is a member of the IL-1 family of proteins. It is a cytokine that binds to and signals through the IL1RL2/IL-36R receptor which in turn activates NF-kappa-B and MAPK signaling pathways in target cells linked to a pro-inflammatory response. IL-36B is synthesized in several cells including resting and activated monocytes, and B cells. The receptor for IL-36 beta is thought to be a combination of IL-1 Rrp2 and IL-1 RAcP. Interleukin 36 beta is one part of the IL-36 signaling system that is thought to be present in epithelial barriers and to take part in local inflammatory response; similar to the IL-1 system with which it shares the coreceptor IL1RAP. Interleukin 36 beta are involved in a number of fundamental biological processes such as stimulating production of interleukin-6 and interleukin-8 in synovial fibroblasts, articular chondrocytes and mature adipocytes, inducing expression of a number of antimicrobial peptides including beta-defensin 4 and beta-defensin 103 as well as a number of matrix metalloproteases, inducing the production of proinflammatory cytokines in bone marrow-derived dendritic cells (BMDCs), including IL-12, IL-1 beta, IL-6, TNF-alpha and IL-23, and activating p38 MAPK phosphorylation in BMDCs. Moreover, interleukin 36 beta may be involved in skin inflammatory response by acting on keratinocytes, dendritic cells, and indirectly on T cells to drive tissue infiltration, cell maturation and cell proliferation. It plays an important role in dendritic cell maturation by stimulating the surface expression of CD80, CD86 and MHC class II and inducing the production of IFN-gamma, IL-4 and IL-17 by T helper 1 (Th1) cells, cultured CD4+ T cells and splenocytes.

Product Name: Recombinant Mouse IL-36b
Catalog #: PEM0926



注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.