


## 概述 (Summary)

|                             |   |
|-----------------------------|---|
| 英文全称                        | CCL5/C-C Motif Chemokine 5/RANTES   |
| 纯度 (Purity)                 | Greater than 95% as determined by reducing SDS-PAGE   |
| 内毒素 (Endotoxin level)       | <1 EU/μg as determined by LAL test.   |
| 蛋白构建 (Construction)         | Recombinant Rat C-C Motif Chemokine 5 is produced by our E.coli expression system and the target gene encoding Ser25-Ser92 is expressed with a 6His tag at the N-terminus.  |
| Accession #                 | P50231  |
| 蛋白标签 (Tag)                  |   |
| 表达宿主 (Host)                 | E.coli  |
| 种属 (Species)                | Rat   |
| 预测分子量 (Predicted MW)        | 10 KDa  |
| 蛋白形态 (Form)                 | Lyophilized from a 0.2 μm filtered solution of 20mM PB, 500mM NaCl, 2mM EDTA, pH 7.4 .  |
| 储存缓冲液 (Buffer)              |   |
| 运输方式 (Shipping)             | The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.  |
| 稳定性&储存 (Stability &Storage) | Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.   |
| 复溶 (Reconstitution)         | Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. |

## 电泳图 (SDS-PAGE image)



## 背景 (Background)

### 分子別名 (Alternative Names)

C-C motif chemokine 5; SIS-delta; Small-inducible cytokine A5; T-cell-specific protein RANTES; Ccl5; Scya5

### 背景介绍 (References)

C-C motif chemokine 5(CCL5) is a  $\beta$ -chemokine that plays a primary role in the inflammatory immune response by means of its ability to attract and activate leukocytes. CCL5 is secreted by many cell types at inflammatory sites, and it exerts a wide range of activities through the receptors CCR1, CCR3, CCR4, and CCR5. Inflammatory responses can be impaired by the sequestration of CCL5 by the cytomegalovirus protein US28. Oligomerization of CCL5 on glycosaminoglycans is required for CCR1mediated leukocyte adhesion and activation as well as CCL5' s interaction with the chemokine CXCL4/PF4. The deposition of CCL5 on activated vascular endothelial cells is crucial for monocyte adhesion to damaged vasculature, but CCL5 oligomerization is not required for the extravasation of adherent leukocytes. CCL5 is upregulated in breast cancer and promotes tumor progression through the attraction of proinflammatory macrophages in addition to its actions on tumor cells, stromal cells, and the vasculature.

## 注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.