


概述 (Summary)

英文全称	TSLP/Thymic stromal lymphopoietin
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Cynomolgus Thymic Stromal Lymphopoietin is produced by our E.coli expression system and the target gene encoding Tyr29-Gln159(Glu37Gln) is expressed with a 6His tag at the C-terminus.
Accession #	XP_005557555.1
蛋白标签 (Tag)	
表达宿主 (Host)	E.coli
种属 (Species)	Cynomolgus
预测分子量 (Predicted MW)	16.2 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of 20mM Tris-HCl, 6% Trehalose, 2% Glycine, 50mM NaCl, 0.05% Tween 80, pH7.5.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Cynomolgus TSLP (C-6His)
Catalog #: PEV2124



背景 (Background)

分子别名 (Alternative Names)

Thymic stromal lymphopoietin;Thymic stroma-derived lymphopoietin;Tslp

背景介绍 (References)

Thymic stromal lymphopoietin (TSLP) is a protein belonging to the cytokine family, contains 140 amino acids. It is known to play an important role in the maturation of T cell populations through activation of antigen presenting cells. TSLP induces the release of T-cell-attracting chemokines from monocytes and, in particular, enhances the maturation of CD11c+ dendritic cells. It can induce allergic inflammation by directly activating mast cells. TSLP is produced mainly by non-hematopoietic cells such as fibroblasts, epithelial cells and different types of stromal or stromal-like cells. These cells are located in regions where TSLP activity is required.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.