Product Name: Recombinant Human AGP2 (C-6His)

Catalog #: PHH0046



概述 (Summary)

英文全称 alpha-1-acid glycoprotein 2/AGP2/ORM2

纯度 (Purity) Greater than 95% as determined by reducing SDS-PAGE

内毒素 (Endotoxin level) <1 EU/μg as determined by LAL test.

蛋白构建 (Construction) Recombinant Human Alpha-1-acid Glycoprotein 2 is produced by our

Mammalian expression system and the target gene encoding Gln19-

Ser201 is expressed with a 6His tag at the C-terminus.

Accession # P19652

蛋白标签 (Tag)

表达宿主 (Host) Human Cells

种属 (Species)Human预测分子量 (Predicted MW)22.6 KDa

蛋白形态 (Form) Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.

储存缓冲液 (Buffer)

运输方式 (Shipping) The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

稳定性&储存 (Stability &Storage) Lyophilized protein should be stored at ≤ -20°C, stable for one year after

receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days.

Aliquots of reconstituted samples are stable at \leq -20°C for 3 months.

复溶 (Reconstitution) Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It

Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to

is not recommended to reconstitute to a concentration less than 100µg/ml.

reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize

freeze-thaw cycles.


电泳图 (SDS-PAGE image)

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838

Product Name: Recombinant Human AGP2 (C-6His)

Catalog #: PHH0046





背景 (Background)

分子別名 (Alternative Names) 背景介绍 (References) Orosomucoid-2; OMD 2; ORM2; AGP2

Alpha-1-acid glycoprotein 2 is a secreted protein that belongs to the calycin superfamily, lipocalin family. It is expressed by the liver and secreted in plasma. It appears to function in modulating the activity of the immune system during the acute-phase reaction. It functions as transport protein in the blood stream. It binds various hydrophobic ligands in the interior of its beta-barrel domain. It also binds synthetic drugs and influences their distribution and availability.

注意事项 (Note)

For Research Use Only, Not for Diagnostic Use.