


概述 (Summary)

英文全称	Apolipoprotein A-II/ApoA2
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Human Apolipoprotein A-II is produced by our Mammalian expression system and the target gene encoding Gln24-Gln100 is expressed with a 6His tag at the C-terminus.
Accession #	P02652
蛋白标签 (Tag)	
表达宿主 (Host)	Human Cells
种属 (Species)	Human
预测分子量 (Predicted MW)	9.74 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Human ApoA2 (C-6His)
Catalog #: PHH0083



背景 (Background)

分子别名 (Alternative Names)

Apolipoprotein A-II;Apo-AII;Apolipoprotein A2;Truncated apolipoprotein A-II;ProapoA-II;APOA2

背景介绍 (References)

Apolipoprotein A-II(Apo-AII for short), also known as Apolipoprotein A2, is a secreted protein which belongs to the apolipoprotein A2 family. It exists as a disulfide-linked homodimer; and also can form a disulfide-linked heterodimer with APOD. APOA2 is the 2nd most abundant protein of the high density lipoprotein particles. This protein may stabilize HDL (high density lipoprotein) structure by its association with lipids, and affect the HDL metabolism. Defects in APOA2 gene might cause apolipoprotein A-II deficiency or hypercholesterolemia.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.