


概述 (Summary)

英文全称	Asialoglycoprotein receptor 1/ASGPR1
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Human Asialoglycoprotein Receptor 1 is produced by our Mammalian expression system and the target gene encoding Gln62-Ile291 is expressed with a 6His tag at the C-terminus.
Accession #	P07306
蛋白标签 (Tag)	
表达宿主 (Host)	Human Cells
种属 (Species)	Human
预测分子量 (Predicted MW)	27.43 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.2.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Human ASGPR1 (C-6His)
Catalog #: PHH0105



背景 (Background)

分子别名 (Alternative Names)

Asialoglycoprotein Receptor 1; ASGP-R 1; ASGPR 1; C-Type Lectin Domain Family 4 Member H1; Hepatic Lectin H1; HL-1; ASGR1; CLEC4H1

背景介绍 (References)

Asialoglycoprotein Receptor 1 (ASGPR1) is an endocytic recycling receptor, belongs to the long-form subfamily of the C-type/Ca²⁺-dependent lectin family. ASGPR consists of two noncovalently-linked subunits, ASGPR1 and ASGPR2. ASGPR1 mediates the endocytosis of plasma glycoproteins, recognizes terminal galactose and N-acetylgalactosamine units. When the ligand binds to ASGPR1, results in the complex is internalized and transported to a sorting organelle, then ASGPR1 and ligand can be disassociated, ASGPR1 returns to the cell membrane surface.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.