产品名称：Recombinant Human BCMA (C-Fc)
产品货号：PHH0140

概述（Summary）

英文全称	BCMA/TNFRSF17/CD269/B cell maturation antigen
纯度（Purity）	Greater than 95% as determined by reducing SDS-PAGE
内毒素（Endotoxin level）	<1 EU/μg as determined by LAL test.
蛋白构建（Construction）	Recombinant Human B-cell Maturation Protein is produced by our Mammalian expression system and the target gene encoding Met1-Ala54 is expressed with a human IgG1 Fc tag at the C-terminus.
Accession #	Q02223
表达宿主（Host）	Human Cells
种属（Species）	Human
预测分子量（Predicted MW）	32.8 KDa
制剂（Form）	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
运输方式（Shipping）	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存（Stability &Storage）	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
复溶（Reconstitution）	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图（SDS-PAGE image）

产品名称：Recombinant Human BCMA (C-Fc)
产品货号：PHH0140



背景（Background）

分子别名（Alternative Names）	Tumor necrosis factor receptor superfamily member 17;TNFRSF17;B-cell maturation protein;CD269
背景介绍（References）	B cell maturation protein (BCMA) is a type III membrane protein which belongs to the TNF receptor superfamily (TNFRSF17). BCMA contains one extracellular cysteine rich domain, as well as TACI. Human BCMA is a 184 amino acid (aa) protein consisting of a 54 aa extracellular domain, a 23 aa transmembrane domain, and a 107 aa intracellular domain. Mouse and human BCMA share 62% amino acid identity. BCMA is mainly expressed in immune organs and mature B cell lines. BCMA appears to be localized to the Golgi compartment. BCMA may play an important role in B cell development, function and regulation.

注意事项（Note）

For research use only .