Product Name: Recombinant Human BOC (C-6His)

Catalog #: PHH0163



概述 (Summary)

英文全称 BOC protein/BOC

纯度 (Purity) Greater than 95% as determined by reducing SDS-PAGE

内毒素 (Endotoxin level) <1 EU/μg as determined by LAL test.

蛋白构建 (Construction) Recombinant Human Brother of CDO is produced by our Mammalian

expression system and the target gene encoding Asp31-Ser157 is

expressed with a 6His tag at the C-terminus.

Accession # Q96DN7

蛋白标签 (Tag)

表达宿主 (Host) Human Cells

种属 (Species)Human预测分子量 (Predicted MW)14.2 KDa

蛋白形态 (Form) Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.

储存缓冲液 (Buffer)

运输方式 (Shipping) The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

稳定性&储存 (Stability &Storage) Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3

months under sterile conditions after opening. Please minimize freeze-thaw

cycles.

复溶 (Reconstitution) Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It

is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized

protein in distilled water. Please aliquot the reconstituted solution to minimize

freeze-thaw cycles.


电泳图 (SDS-PAGE image)

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838

Product Name: Recombinant Human BOC (C-6His)

Catalog #: PHH0163





背景 (Background)

分子别名 (Alternative Names) 背景介绍 (References) BOC protein; Boc homolog (Mouse); isoform CRA_b; Brother of CDO Brother of CDO is a protein that in humans is encoded by the BOC gene. CDON and BOC are cell surface receptors of the immunoglobulin (Ig)/fibronectin type III repeat family involved in myogenic differentiation. CDON and BOC are coexpressed during development, form complexes with each other in a cis fashion, and are related to each other in their ectodomains, but each has a unique long cytoplasmic tail.

注意事项 (Note)

For Research Use Only, Not for Diagnostic Use.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838