

Product Name: Recombinant Human BTN3A2 (C-6His)
Catalog #: PHH0178




概述 (Summary)

英文全称	BTN3A2/Butyrophilin subfamily 3 member A2
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Human Butyrophilin Subfamily 3 Member A2 is produced by our Mammalian expression system and the target gene encoding Gln30-Trp248 is expressed with a 6His tag at the C-terminus.
Accession #	P78410
蛋白标签 (Tag)	
表达宿主 (Host)	Human Cells
种属 (Species)	Human
预测分子量 (Predicted MW)	24.6 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Human BTN3A2 (C-6His)
Catalog #: PHH0178



背景 (Background)

分子别名 (Alternative Names)

Butyrophilin subfamily 3 member A2; BT3.2; BTF3; BTF4;BTN3A2

背景介绍 (References)

Butyrophilin subfamily 3 member A2, also known as BT3.2, BTF3, BTF4 and BTN3A2, is a single-pass type I membrane protein. It is a member of the butyrophilin (BTN) family and the immunoglobulin (Ig) superfamily. Mature human BTN3A2 is a 305 amino acid (aa) glycoprotein. It contains a 219 aa extracellular region with one V-type Ig-like domain, and a 65 aa cytoplasmic tail. The cytoplasmic region undergoes phosphorylation on two serines. There are three potential splice forms. BTN3A2 is postulated to be expressed on immune-related cells, as it has a structural similarity to MHC and CD80/CD86 molecules. It plays a role in T-cell responses in the adaptive immune response and inhibits the release of IFNG from activated T-cells.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.