


概述 (Summary)

英文全称	Cadherin-17/CDH17
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Human Cadherin-17 is produced by our Mammalian expression system and the target gene encoding Gln23-Met787 is expressed with a 6His tag at the C-terminus.
Accession #	AAI13465.1
蛋白标签 (Tag)	
表达宿主 (Host)	Human Cells
种属 (Species)	Human
预测分子量 (Predicted MW)	86 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Human CDH17 (C-6His)
Catalog #: PHH0192



背景 (Background)

分子别名 (Alternative Names)

Cadherin-17; Intestinal Peptide-Associated Transporter HPT-1; Liver-Intestine Cadherin; LI-Cadherin; CDH17

背景介绍 (References)

Cadherin-17 is a single-pass type I membrane protein that belongs to the cadherin superfamily. Cadherin-17 consists of one extracellular region containing seven cadherin domains and one transmembrane region but it lacks the conserved cytoplasmic domain. Cadherin-17 is expressed in the gastrointestinal tract and pancreatic duct. Cadherins are calcium dependent cell adhesion proteins and preferentially interact with each other in a homophilic manner in connecting cells. Cadherin-17 may have a role in the morphological organization of liver and intestine and involved in intestinal peptide transport.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.