


概述 (Summary)

英文全称	CXADR/CAR/NR1I3
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Human Cocksackievirus And Adenovirus Receptor is produced by our Mammalian expression system and the target gene encoding Leu20-Gly237 is expressed with a 6His tag at the C-terminus.
Accession #	P78310
蛋白标签 (Tag)	
表达宿主 (Host)	Human Cells
种属 (Species)	Human
预测分子量 (Predicted MW)	25.08 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.2.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Human CAR (C-6His)
Catalog #: PHH0209



背景 (Background)

分子别名 (Alternative Names)

Coxsackievirus and Adenovirus Receptor; CAR; hCAR; CVB3-Binding Protein; Coxsackievirus B-Adenovirus Receptor; HCVADR; CXADR; CAR

背景介绍 (References)

Coxsackievirus and Adenovirus Receptor (CAR) belongs to the CTX family of the Ig superfamily. CXADR is a type I transmembrane glycoprotein and expressed in pancreas, brain, heart, small intestine, testis, prostate. It is a receptor that mediates gene transfer and also act as an adhesion molecule within junctional complexes, notably between epithelial cells lining body cavities and within myocardial intercalated discs. CXADR contains an extracellular domain, a transmembrane helix and a C-terminal intracellular domain. The C-terminal interacts with few cytoplasmic junctional proteins, microtubules and the actin cytoskeleton.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.