


概述 (Summary)

英文全称	CCL24/Eotaxin-2/MPIF-2/C-C motif chemokine 24
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Human C-C Motif Chemokine 24 is produced by our Mammalian expression system and the target gene encoding Val27-Cys119 is expressed with a 6His tag at the C-terminus.
Accession #	O00175
蛋白标签 (Tag)	
表达宿主 (Host)	Human Cells
种属 (Species)	Human
预测分子量 (Predicted MW)	11.5 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)



背景 (Background)

分子别名 (Alternative Names)

C-C motif chemokine 24 (CCL24); CK-beta-6; Eosinophil chemotactic protein; Eotaxin-2; Myeloid progenitor inhibitory factor 2; Small-inducible cytokine A24; Ckb-6; MPIF-2; SCYA24

背景介绍 (References)

Cytokine are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The CC cytokines are proteins characterized by two adjacent cysteine. CCL24 displays the chemotactic for resting T-lymphocytes and eosinophils. CCL24 has lower chemotactic activity for neutrophils, none for monocytes and activated lymphocytes. CCL24 is a strong suppressor of colony formation by a multipotential hematopoietic progenitor cell lines.

注意事项 (Note)

For Research Use Only, Not for Diagnostic Use.