


概述 (Summary)

英文全称	CD113/Nectin-3/PVRL3
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Human Poliovirus Receptor-Related Protein 3 is produced by our Mammalian expression system and the target gene encoding Gly58-Cys366 is expressed with a 6His tag at the C-terminus.
Accession #	Q9NQS3
蛋白标签 (Tag)	
表达宿主 (Host)	Human Cells
种属 (Species)	Human
预测分子量 (Predicted MW)	34.96 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, 5% Trehalose, pH 7.4.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Human Nectin-3 (C-6His)
Catalog #: PHH0284



背景 (Background)

分子别名 (Alternative Names)

Poliovirus Receptor-Related Protein 3; CDw113; Nectin-3; CD113; PVRL3; PRR3

背景介绍 (References)

Nectin-3 is a type I transmembrane glycoprotein that belongs to the nectin family. Its precursor is 549 amino acids in length and contains an extended signal sequence of 57 amino acids, an extracellular domain (ECD) of 347 amino acids, a transmembrane segment of 21 amino acids, and a cytoplasmic region of 124 amino acids. It is predominantly expressed in testis and placenta as well as in various cell lines, including epithelial cell lines. Nectin-3 plays a role in cell-cell adhesion through heterophilic trans-interactions with nectin-like proteins or nectins, such as trans-interaction with PVRL2/Nectin-2 at Sertoli-spermatid junctions. Nectin-3 is also involved in the formation of cell-cell junctions, including adherens junctions and synapses.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.