


概述 (Summary)

英文全称	Syndecan-1/SDC1/CD138
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Human Syndecan-1 is produced by our Mammalian expression system and the target gene encoding Gln18-Glu251 is expressed with a 6His tag at the C-terminus.
Accession #	AAH08765.1
蛋白标签 (Tag)	
表达宿主 (Host)	Human Cells
种属 (Species)	Human
预测分子量 (Predicted MW)	25.2 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Human SDC1 (C-6His)
Catalog #: PHH0290



背景 (Background)

分子别名 (Alternative Names)

Syndecan-1; SYND1; CD138; SDC1; SDC

背景介绍 (References)

Syndecan-1 is a single-pass type I membrane protein that belongs to the syndecan proteoglycan family. The syndecans mediate cell binding, cell signaling, and cytoskeletal organization and syndecan receptors are required for internalization of the HIV-1 tat protein. Human SDC1 is synthesized as a 310 amino acid precursor that contains a 22 amino acid signal sequence, and a 288 amino acid mature chain. The Syndecan-1 protein functions as an integral membrane protein and participates in cell proliferation, cell migration and cell-matrix interactions via its receptor for extracellular matrix proteins. Altered Syndecan-1 expression has been detected in several different tumor types.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.