


概述 (Summary)

英文全称	ALCAM/CD166
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Human CD166 Antigen is produced by our Mammalian expression system and the target gene encoding Trp28-Ala526 is expressed with a human IgG1 Fc tag at the C-terminus.
Accession #	Q13740
蛋白标签 (Tag)	
表达宿主 (Host)	Human Cells
种属 (Species)	Human
预测分子量 (Predicted MW)	82.7 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Human ALCAM (C-Fc)
Catalog #: PHH0300



背景 (Background)

分子别名 (Alternative Names)

CD166 antigen; Activated leukocyte cell adhesion molecule; CD166; ALCAM; MEMD

背景介绍 (References)

Activated leukocyte cell adhesion molecule (ALCAM), also named as CD166 and MEMD, is a type I transmembrane glycoprotein of immunoglobulin superfamily, which mediates homotypic and heterotypic interactions between cells. ALCAM is expressed on thymic epithelium, microvascular endothelium, activated lymphocytes and monocytes, and monocytederived dendritic cells. ALCAM mediates low-affinity adhesion with itself or the cysteine-rich scavenger receptor CD6 to regulate T cell development, immunological synapses (IS), and cell migration through endothelial junctions. ALCAM on thymic epithelia mediates adhesion to CD6 on CD4+CD8+ T cells. Adhesion of ALCAM expressing antigen presenting cells and CD6-expressing T cells stabilizes the early IS, while later it enhances CD3 effects on T cell proliferation, CD25 expression, and Th1 commitment. ALCAM may influence expression or adhesion of the neuronal adhesion molecule NCAM1, both in the developing retina and invasive melanoma.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.