


概述 (Summary)

英文全称	CD177/PRV1
纯度 (Purity)	Greater than 95% as determined by reducing SDS-PAGE
内毒素 (Endotoxin level)	<1 EU/μg as determined by LAL test.
蛋白构建 (Construction)	Recombinant Human Polycythemia Rubra Vera Protein 1 is produced by our Mammalian expression system and the target gene encoding Leu22-Gly407 is expressed with a 6His tag at the C-terminus.
Accession #	AAH29167.1
蛋白标签 (Tag)	
表达宿主 (Host)	Human Cells
种属 (Species)	Human
预测分子量 (Predicted MW)	42.33 KDa
蛋白形态 (Form)	Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.2.
储存缓冲液 (Buffer)	
运输方式 (Shipping)	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存 (Stability &Storage)	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
复溶 (Reconstitution)	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图 (SDS-PAGE image)

Product Name: Recombinant Human CD177 (C-6His)
Catalog #: PHH0304



背景 (Background)

分子别名 (Alternative Names)

CD177 Antigen; Human Neutrophil Alloantigen 2a; HNA-2a; NB1 Glycoprotein; NB1 GP; Polycythemia Rubra Vera Protein 1; PRV-1; CD177; NB1; PRV1

背景介绍 (References)

CD177 is polymorphic and has at least two alleles: PRV1 and NB1. Human PRV1 is a Glycosyl-Phosphatidylinositol (GPI)-linked cell surface glycoprotein that belongs to the uPAR/CD59/Ly6 family of receptors. PRV1 is expressed by neutrophils and neutrophil precursors, and changes in expression serve as diagnostic markers for myeloproliferative disorders such as polycythemia vera and essential thrombocythemia. PRV1 may also be expressed by Erythroblasts, B cells, and Monocytes. NB1, a Glycosyl-Phosphatidylinositol (GPI)-linked cell surface glycoprotein, was first described in a case of neonatal alloimmune neutropenia. It is reported that CD177 functions as a novel heterophilic binding partner that engages PECAM-1 in membrane-proximal IgD6.

注意事项 (Note)

For Research Use Only , Not for Diagnostic Use.